



King's Research Portal

DOI:

[10.1111/jgs.14106](https://doi.org/10.1111/jgs.14106)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Hazra, N. C., Dregan, A., Jackson, S., & Gulliford, M. C. (2016). Drug utilization and inappropriate prescribing in centenarians. *Journal of the American Geriatrics Society*, 64(5), 1079–1084 . <https://doi.org/10.1111/jgs.14106>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Drug Utilization and Inappropriate Prescribing in Centenarians

Nisha C. Hazra, MSc,* Alex Dregan, PhD,*[†] Stephen Jackson, MD,[‡] and Martin C. Gulliford, FPPH*[†]

OBJECTIVES: To use primary care electronic health records (EHRs) to evaluate prescriptions and inappropriate prescribing in men and women at age 100.

DESIGN: Population-based cohort study.

SETTING: Primary care database in the United Kingdom, 1990 to 2013.

PARTICIPANTS: Individuals reaching the age of 100 between 1990 and 2013 (N = 11,084; n = 8,982 women, n = 2,102 men).

MEASUREMENTS: Main drug classes prescribed and potentially inappropriate prescribing according to the 2012 American Geriatrics Society Beers Criteria.

RESULTS: At the age of 100, 73% of individuals (79% of women, 54% of men) had received one or more prescription drugs, with a median of 7 (interquartile range 0–12) prescription items. The most frequently prescribed drug classes were cardiovascular (53%), central nervous system (CNS) (53%), and gastrointestinal (47%). Overall, 32% of participants (28% of men, 32% of women) who received drug prescriptions may have received one or more potentially inappropriate prescriptions, with temazepam and amitriptyline being the most frequent. CNS prescriptions were potentially inappropriate in 23% of individuals, and anticholinergic prescriptions were potentially inappropriate in 18% of individuals.

CONCLUSION: The majority of centenarians are prescribed one or more drug therapies, and the prescription may be inappropriate for up to one-third of these individuals. Research using EHRs offers opportunities to understand prescribing trends and improve pharmacological care of the oldest adults. *J Am Geriatr Soc* 2016.

Key words: centenarians; epidemiology; inappropriate prescribing; aging; primary care

Remarkable reductions in mortality are associated with rapid increases in life expectancy in many developed countries. Recent estimates in the United Kingdom (UK) indicate a quintupling of the number of centenarians over the past 3 decades, to 13,780 in 2013.¹ This trend is expected to continue, with approximately 68,000 centenarians in the United Kingdom² and 1 million centenarians worldwide by 2030.^{3,4} Greater longevity is likely to be associated with higher rates of drug prescribing and greater expenditures on health services.⁵ Elderly people consume 45% to 55% of all drugs in the United Kingdom,^{6,7} and 38% of all drugs in the United States,⁸ but few studies in the United Kingdom have described drug utilization and none have described potentially inappropriate prescribing (PIP) in the oldest adults.

Drug prescribing and managing medication use in older adults is complex and challenging because multiple morbidity, frailty, and cognitive impairment often accompany old age. Drug prescribing in very old adults is challenging to clinicians because very old adults often require multiple drug therapy but are also at great risk of drug-related adverse effects. Evidence regarding drug prescribing in centenarians is rare and, when available, rather inconsistent.^{9–12} Existing studies include self-reported data and tend to focus on younger old people,^{7,9,10,12} often excluding centenarians.^{9,13,14}

Electronic health records (EHRs) are a valuable resource to explore the epidemiology of drug prescribing in population-based samples.¹² Electronic data can help overcome biases in self-reported prescribing patterns and medical histories, a common limitation of previous studies.^{7,15–17} The central role of general practice within the U.K. National Health Service allows for long-term prescribing to be managed in primary care. This study aimed to use primary care EHR data from the U.K. Clinical Practice Research Datalink (CPRD), one of the world's largest primary care databases, to describe drug use and to estimate the frequency of PIP in a large cohort of centenarians.

From the *Department of Primary Care and Public Health Sciences, King's College London; [†]Biomedical Research Centre, National Institute for Health Research, Guy's and St Thomas' National Health Service Foundation Trust; and [‡]Department of Clinical Gerontology, King's College Hospital, London, UK.

Address correspondence to Nisha C. Hazra, Department of Primary Care and Public Health Sciences, King's College London, 7th Floor Capital House, 42 Weston St, London SE1 3QD, UK. E-mail: nisha.hazra@kcl.ac.uk

DOI: 10.1111/jgs.14106

METHODS

Setting and Study Design

The study included a nationally representative sample of individuals in primary care from the CPRD who reached the age of 100 between 1990 and 2013, as described previously.¹⁸ Practices can contribute data only when it is deemed of sufficient quality for research, according to a set of criteria that the CPRD group has developed.¹⁹ This study, which received scientific and ethical approval from the Independent Scientific Advisory Committee for CPRD studies (ISAC Protocol 13_151), is based on fully anonymized data, and research ethics committee approval was not required.

Participants

A population-based cohort of centenarians was drawn from the CPRD between January 1, 1990, and September 30, 2013, as previously described.¹⁸ Eligible participants were registered at CPRD general practices during the year in which they turned 100.

Measures

Drug utilization was first evaluated using the main chapters from the British National Formulary (BNF) as categories. The BNF is published every 6 months and provides healthcare professionals with up-to-date information about medicines.²⁰ Each participant's record was evaluated for drug prescriptions in the following drug categories: gastrointestinal; cardiovascular; respiratory; central nervous system (CNS); antimicrobial; endocrine; gynecological and urinary tract; neoplasms and immunosuppression; nutrition and blood; musculoskeletal; eye; ear, nose, and oropharynx; skin; immunological; and anesthetic. Centenarians were considered to have been prescribed a drug from a particular category if at least one relevant drug was prescribed during their 100th year. Drugs relevant to more than one category were counted in each. For example, prednisolone is included in the endocrine and gastrointestinal categories. Descriptive statistics were used to estimate the total number of prescriptions before reaching the age of 100, from the 95th year to the 100th year of life.

The frequency of PIP among centenarians was evaluated using the updated 2012 American Geriatrics Society (AGS) Beers Criteria.²¹ An expert group developed the criteria in 1991,²² with the latest revision by the AGS in 2012,²¹ to provide explicit criteria for PIP in elderly adults. The criteria include three primary groups: a list of potentially inappropriate medication (PIMs) to avoid independent of disease or condition; a list of medications to be avoided in older persons with specific diseases or conditions; and a list of PIMs to be used with caution in older adults.²¹ This tool has been widely used to evaluate PIP of drugs to elderly individuals in a variety of settings.^{15–17,23–25}

In the present study, we estimated the proportion of centenarians receiving one or more prescriptions from each overall Beers category, independent of disease or condition, including—anticholinergics, anti-thrombotics, anti-infectives, cardiovascular, CNS, endocrine, gastrointestinal, and analgesics. The proportion prescribed at least one specific

PIM within each category was also estimated. The standard criteria were used, rather than the disease-specific criteria, because of the large population-based sample in the study. The three drugs that were most frequently prescribed inappropriately were evaluated, and descriptive statistics were used to determine the median number of PIMs in each category. Desiccated thyroid is not listed in the BNF, so it was excluded from the analysis. Glyburide is known as glibenclamide in the United Kingdom. The prescribing of nonsteroidal anti-inflammatory drugs (NSAIDs) in this study was considered inappropriate only if, as stated in the 2012 AGS Beers Criteria, they were not taken with a gastroprotective agent. All topical non-cyclooxygenase-selective NSAIDs were not considered as inappropriate and only oral formulations were included, as stated by the 2012 AGS Beers Criteria. Stata version 13.0 was used to conduct all analysis (Stata Corp., College Station, TX).

RESULTS

A cohort of 11,047 centenarians (8,982 women, 2,102 men), who reached the age of 100 between 1990 and 2013, was selected for analysis. Eighty-four percent were born between 1900 and 1913 and the remaining 16% between 1890 and 1899. The median annual number of prescriptions was 7 (interquartile range 0–12) during the 100th year.

Table 1 shows the most frequently prescribed drugs at the age of 100. Drug utilization for each drug class was higher in women than in men, except for urinary tract

Table 1. Frequency of Different Categories of Prescriptions According to Sex

Drug Class from British National Formulary	Female, n = 8,982	Male, n = 2,102	All, N = 11,084
	n (%) ^a		
Any	6,904 (79)	1,136 (54)	8,040 (73)
Cardiovascular	5,044 (56)	835 (40)	5,879 (53)
Central nervous system ^b	5,182 (58)	741 (35)	5,923 (53)
Gastrointestinal	4,488 (50)	752 (36)	5,240 (47)
Skin	3,524 (39)	537 (26)	4,061 (37)
Nutrition and blood	3,485 (39)	479 (23)	3,964 (36)
Antimicrobial	4,028 (45)	631 (30)	4,659 (42)
Eye	2,038 (23)	316 (15)	2,354 (21)
Musculoskeletal and joint diseases	1,963 (22)	307 (15)	2,270 (20)
Endocrine	1,818 (20)	283 (13)	2,101 (19)
Respiratory	1,545 (17)	277 (13)	1,822 (16)
Anesthesia	1,131 (13)	198 (9)	1,329 (12)
Ear, nose, oropharynx	828 (9)	171 (8)	999 (9)
Immunological products and vaccines	786 (9)	119 (6)	905 (8)
Gynecological and urinary tract	692 (8)	206 (10)	898 (8)
Neoplasms and immunosuppression	497 (6)	92 (4)	589 (5)

^aFrequency of individuals with at least one prescription during the year they turned 100.

^bHypnotics, anxiolytics, antidepressants, analgesics, drugs for nausea and vertigo, antiepileptics, drugs for parkinsonism and dementia.

drugs. The most frequently prescribed drugs overall were those affecting the CNS (53%), including hypnotics, anxiolytics, antidepressants, analgesics, drugs for nausea and vertigo, antiepileptics, and antidementia, as well as cardiovascular (53%) and gastrointestinal (47%) drugs. The distribution of prescriptions issued to men and women centenarians is shown in Figure 1. More than twice as many men as women did not receive any prescriptions in their 100th year, and the proportion of women was higher than men for each increasing category of multiple prescriptions.

The frequency of PIP according to the 2012 AGS Beers Criteria is presented in Table 2. Overall, 32% of centenarians were prescribed a PIM, and the three most frequently prescribed PIMs were temazepam, amitriptyline, and nitrofurantoin. The drug class with the highest proportion of PIP was CNS medications (23%). Approximately one-fifth of centenarians (19% women, 13% men) received anticholinergic drugs, with PIMs including chlorphenamine, hydroxyzine, and promethazine hydrochloride. Despite the high levels of prescribing for gastrointestinal, cardiovascular, and analgesic drugs, low levels of inappropriate prescribing (5%) were observed within these classes.

DISCUSSION

To the knowledge of the authors, this is the first population-based study describing medication use and PIP in centenarians. Only a minority of centenarians did not receive prescription medicines, with a higher proportion of men not receiving any prescriptions, consistent with their superior health status at age 100.¹⁸ Almost 80% of women and 54% of men were prescribed at least one drug during their 100th year. Sex differences in overall prescribing between centenarians have not been reported previously. This disparity could be because of sex differences in health-seeking behavior, for example, not seeking a physician's advice, or nonadherence, as well as resulting from differences in health status. Up to one-third of centenarians were prescribed a PIM. The highest frequencies of PIP were attributed to the use of benzodiazepines (temazepam, diazepam), amitriptyline, and nitrofurantoin. The 2012 AGS Beers Criteria recommendation to avoid all of these drugs is

“strong,” and the reported quality of evidence is “high” for avoiding temazepam, diazepam, and amitriptyline and “moderate” for avoiding nitrofurantoin.²¹ The 2012 AGS Beers Criteria recommend avoiding nitrofurantoin in individuals with creatinine clearance less than 60 mL/min because of concerns about lack of efficacy from inadequate drug concentrations in the urine. In view of the advanced age of the cohort, it is likely that most will have poor renal function and should therefore be using safer alternatives such as ciprofloxacin or trimethoprim.²⁶

Comparison with Existing Literature

Existing studies on drug utilization in elderly adults have often focused on younger cohorts of old people^{7,9,10,12} or do not include centenarians.^{9,13,14} There is scarce evidence about PIP in extreme old age. These studies focusing on younger elderly adults tend to rely on self-reported questionnaires and interviews,^{7,15–17,24,27} resulting in a high risk of responder bias.

There have been no previous studies reporting PIP in a large group of centenarians. Reports of prevalence of PIP in individuals aged 65 and older are inconsistent, and few studies have used the updated 2012 criteria. One study²⁷ using the 2012 criteria reported a higher prevalence of PIP (44%) in Spanish individuals aged 65 and older than the present findings (32%) in centenarians. The most frequently prescribed PIMs were benzodiazepines, similar to the present data. Another study in New Zealand²⁴ also reported a higher PIM prevalence (42.7%) for community-dwelling individuals aged 75 and older than for those aged 100 and older in CPRD, whereas another¹⁵ reported a prevalence of 17%. All three studies^{15,24,27} used self-reported data. Several studies reported pain medications as the most commonly prescribed PIMs,^{15,17,23,24} including two studies using the Screening Tool of Older People's potentially inappropriate Prescriptions/Screening Tool to Alert doctors to the Right Treatment (STOPP/START) criteria.^{28,29} This is inconsistent with the present findings in centenarians, reporting CNS medications and anticholinergics as the most commonly prescribed PIMs. It is not stated in all these previous studies whether concurrent use of gastroprotective agents was considered alongside NSAIDs.




Figure 1. Frequency of prescriptions during 100th year according to sex.

Strengths and Limitations

This study had the strengths of a large sample drawn from a representative population of U.K. general practices. In the United Kingdom, approximately 98% of individuals are registered with a family practice, ensuring that the present data are complete and nationally representative. Individuals aged 75 and older have an annual review of medicines,³⁰ and those with four or more medicines are reviewed every 6 months, although this review was not introduced until 2002. Using primary care EHRs allowed for the classification of prescribing according to drug category and of specific PIMs, but data were not available for several variables of interest, including whether an individual lived alone or whether they lived in an urban or rural location. EHRs circumvent the problem of recall bias, a limitation of many drug use studies relying on self-reported questionnaires or interviews to collect data on prescrip-

Table 2. Frequency of Potentially Inappropriate Medications Using 2012 American Geriatrics Society (AGS) Beers Criteria

Drug Category	Female, n = 8,982			Male, n = 2,102			All, N = 11,084		
	Total Drug Class, n ^a	PIM, n (%) ^b	Number of PIMs, Median (IQR)	Total Drug Class, n ^a	PIM, n (%) ^b	Number of PIMs, Median (IQR)	Total Drug Class, n ^a	PIM, n (%) ^b	Number of PIMs, Median (IQR)
Any	6,799	2,208 (32)	5 (2–12)	1,108	309 (28)	4 (1–10)	7,907	2,517 (32)	5 (2–12)
Central nervous system	5,182	1,240 (24)	7 (2–13)	741	147 (20)	5 (2–12)	Temazepam	Temazepam	Temazepam
							Amitriptyline	Dipyrindamole	Amitriptyline
							Nitrofurantoin	Amitriptyline	Nitrofurantoin
Anticholinergic	1,279	237 (19)	2 (1–8)	219	28 (13)	2 (1–6)	5,923	1,387 (23)	6 (2–12)
Anti-infective	4,027	353 (9)	1 (1–2)	631	38 (6)	1 (1–2)	Temazepam	Chlorphenamine	Chlorphenamine
							Amitriptyline	Hydroxyzine	Hydroxyzine
							Diazepam	Promethazine	Promethazine
Analgesic	5,348	258 (5)	2.5 (1–7)	844	31 (4)	1 (1–2)	Chlorphenamine	hydrochloride	hydrochloride
							Nitrofurantoin	Hydroxyzine	Hydroxyzine
							Ibuprofen	Nitrofurantoin	Nitrofurantoin
Gastrointestinal	4,653	238 (5)	1 (1–4)	773	31 (4)	1 (1–2)	Diclofenac	Ibuprofen	Ibuprofen
							Meloxicam	Diclofenac	Diclofenac
							Liquid paraffin (mineral oil)	Meloxicam	Meloxicam
Cardiovascular	5,044	207 (4)	7 (3–13)	835	50 (6)	6 (3–11)	Metoclopramide	Metoclopramide	Metoclopramide
							Amiodarone	Liquid paraffin (mineral oil)	Liquid paraffin (mineral oil)
							Doxazosin	Amiodarone	Amiodarone
Antithrombotic	2,578	87 (3)	6 (2–12)	460	26 (6)	6 (2–12)	Sotalol	Doxazosin	Doxazosin
							Dipyrindamole	Spironolactone	Spironolactone
							Glyburide	Dipyrindamole	Dipyrindamole
Endocrine	1,888	15 (1)	11 (3–13)	311	1 (0.3)	6 (6–6)	Conjugated estrogens	Glyburide	Glyburide
							Megestrol	Conjugated estrogens	Conjugated estrogens
								Megestrol	Megestrol

^aNumber of individuals from total sample with at least one prescription during the year of turning 100 years old in each overall Beers category.

^bPIMs = potentially inappropriate medications within each Beers category as defined by the 2012 AGS Beers Criteria (row percentages).

tions. There is a possibility that some prescriptions will not be filled or consumed, and the data may not capture any over-the-counter or secondary care hospital prescriptions, so the present findings may underestimate drug usage of centenarians. Another limitation is that the sample was exposed to a nonuniform drug formulary because the study considers data over a 23-year span. Medical practice in primary care has evolved over time, and as a result, certain drugs used in 1990 may now be considered inappropriate. Several new drugs have also been introduced that could not be prescribed in 1990.

There are limitations of the 2012 AGS Beers Criteria, given their universal application without careful consideration of an individual's response to each drug.^{15,25} There may also be some drugs used in the United Kingdom that were not captured. In 2003, the STOPP/START criteria were developed to identify potential errors in prescribing and prescribing omission in older people according to physiological system.^{24,28} These criteria were created as a Europe-focused tool but are not as widely used as the 2012 AGS Beers Criteria to evaluate the epidemiology of PIP because of their specificity. Although a more-individualized tool may be favored in clinical practice, this approach is less feasible for epidemiological investigation of population-based samples. The 2012 AGS Beers Criteria were specifically designed to use pharmacy records with minimal additional clinical information so that they could be applied to chart reviews or computerized data sets.²¹

According to the 2012 AGS Beers Criteria, the most commonly prescribed PIMs in the present cohort were benzodiazepines, tertiary tricyclic antidepressants, nitrofurantoin, and ibuprofen. The frequencies of PIP found in this study should be interpreted cautiously because each person's risk:benefit ratio for a drug will depend on his or her physiological and clinical status. Only an individual evaluation of each person will confirm the validity of these interpretations. Nevertheless, population-based studies provide useful epidemiological data on relative frequencies of PIP in large cohorts and help identify the inappropriate medications that are most frequently prescribed. This study also identifies specific drugs that may be given more attention in further research on the determinants of PIP in elderly adults.

Conclusion and Implications for Clinical Practice

Polypharmacy and multimorbidity in elderly adults present health professionals with a significant clinical responsibility. The limited empirically based evidence to guide drug prescription in very old adults means that physicians must base their prescribing decisions on clinical knowledge and prior experience with similar conditions, likely from younger cases. There is an urgent need for studies to explore the efficacy, safety, and harms associated with drug prescribing for different chronic conditions in very old adults. There is also the need to model the adverse clinical and economic consequences of inappropriate therapeutic decision-making in this group.

This is the first study to use primary care EHR data to describe prescribing trends in UK centenarians, as well as the extent of PIP according to drug class. It provides proof of concept for using a large EHR database to evaluate

appropriateness of prescribing and a basis for reevaluating indicators of appropriate prescribing as applied to this age group. This also offers valuable data for the modeling of future healthcare needs and costs of the oldest adults in the United Kingdom.

ACKNOWLEDGMENTS

Conflict of Interest: The study is based in part on data from the CPRD obtained under license from the U.K. Medicines and Healthcare products Regulatory Agency, but the interpretation and conclusions contained in this report are those of the authors alone. There are no conflicts of interest involved. The editor in chief has reviewed the conflict of interest checklist provided by the authors and has determined that the authors have no financial or any other kind of personal conflicts with this paper.

This research was supported by the National Institute for Health Research Biomedical Research Centre based at Guy's and St Thomas' National Health Service Foundation Trust and King's College London. Open Access for this article was funded by King's College London.

Author Contributions: All authors were involved in drafting the article and revising it critically for intellectual content, and all authors approved the final version to be published. Gulliford, Hazra, and Dregan had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Jackson, Gulliford: study conception and design. Hazra, Dregan, Gulliford: acquisition of data. Hazra, Dregan, Jackson, Gulliford: analysis and interpretation of data.

Sponsor's Role: The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

REFERENCES

1. Estimates of the Very Old (including Centenarians), for England and Wales, United Kingdom, 2002 to 2013; 2014. Office for National Statistics [on-line]. Available at http://www.ons.gov.uk/ons/dcp171778_378107.pdf Accessed August 2, 2015.
2. International Data Base, 2013. U.S. Census Bureau [on-line]. Available at <http://www.census.gov/population/international/data/idb/informationGateway.php> Accessed October 24, 2014.
3. Secrets of the Centenarians: Life Begins at 100. 2009. New Scientist [on-line]. Available at http://www.newscientist.com/article/mg20327241.300-secrets-of-the-centenarians-life-begins-at-100.html?full=true#UzF_RIbIaQA Accessed January 29, 2014.
4. World Population Ageing 1950–2050: IV. Demographic Profile of the Older Population; 2011. United Nations DESA [on-line]. Available at <http://www.un.org/esa/population/publications/worldageing19502050/pdf/90chapteriv.pdf> Accessed October 25, 2014.
5. Yang Z, Norton EC, Stearns SC. Longevity and health care expenditures: The real reasons older people spend more. *J Gerontol B Psychol Sci Soc Sci* 2003;58B:S2–S10.
6. Shah RR. Drug development and use in the elderly: Search for the right dose and dosing regimen. *Br J Clin Pharmacol* 2004;58:452–469.
7. Cakir YT, Sonbahar M, Can H et al. Drug usage habits and multiple drug usage of elderly individuals in nursing homes. *Turk J Geriatr* 2014;17:172–179.
8. National Center for Health Statistics Data Brief, No. 41: Population Aging and the Use of Office-based Physician Services. 2010. U.S. Department of Health and Human Services: Centers for Disease Control and Prevention [on-line]. Available at <http://www.cdc.gov/nchs/data/databriefs/db41.pdf> Accessed November 28, 2014.
9. Nobili A, Franchi C, Pasina L et al. Drug utilization and polypharmacy in an Italian elderly population: The EPIFARM-Elderly Project. *Pharmacopidemiol Drug Saf* 2011;20:488–496.

10. Hartikainen S, Klaukka T. Use of psychotropics is high among very old people. *Eur J Clin Pharmacol* 2004;59:849–850.
11. Wastesson JW, Parker MG, Fastbom J et al. Drug use in centenarians compared with nonagenarians and octogenarians in Sweden: A nationwide register-based study. *Age Ageing* 2012;41:218–224.
12. Melzer D, Tavakoly B, Winder RE et al. Much more medicine for the oldest old: Trends in UK electronic clinical records. *Age Ageing* 2015;44:46–53.
13. Kennerfalk A, Ruigomez A, Wallander MA. Geriatric drug therapy and healthcare utilization in the United Kingdom. *Ann Pharmacother* 2002;36:797–803.
14. Razavi M, Meera NK, Karimian H et al. A profile of drug utilization among elderly inpatients admitted at a tertiary level hospital in Bangalore: A prospective study. *Arch Pharma Pract* 2012;3:217–222.
15. Sonmez Y, Asci H, Olgun Izmirli G et al. Evaluation of potentially inappropriate drug use and medical non-adherence in a community-dwelling elderly population: A cross-sectional study. *Turk J Geriatr* 2014;17:125–133.
16. Fialova D, Topinkova E, Gambassi G et al. Potentially inappropriate medication use among elderly home care patients in Europe. *JAMA* 2005;293:1348–1358.
17. Zhan C, Sangl J, Bierman AS et al. Potentially inappropriate medication use in the community-dwelling elderly: Findings from the 1996 Medical Expenditure Panel Survey. *JAMA* 2001;286:2823–2829.
18. Hazra N, Dregan A, Jackson S et al. Gender differences in health at age 100: Population-based cohort study of centenarians using electronic health records. *J Am Geriatr Soc* 2015;63:1331–1338.
19. Independent Scientific Advisory Committee for Medicines and Healthcare products Regulatory Agency (MHRA) Database Research (ISAC): Annual Report. 2012. Clinical Practice Research Datalink [on-line]. Available at http://www.cprd.com/_docs/ISAC%20Annual%20Report_2012.pdf Accessed September 9, 2014.
20. British National Formulary. BMJ Group, RCPCH Publications Ltd and the Royal Pharmaceutical Society of Great Britain 2014 [on-line]. Available at http://www.bnf.org/bnf/org_450002.htm Accessed October 31, 2014.
21. The American Geriatrics Society 2012 Beers Criteria Update Expert Panel. American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc* 2012;60:616–631.
22. Beers MH, Ouslander JG, Rollingher I et al. Explicit criteria for determining inappropriate medication use in nursing homes. *Arch Intern Med* 1991;151:1825–1832.
23. Fahlman C, Lynn J, Finch M et al. Potentially inappropriate medication use by Medicaid+Choice beneficiaries in the last year of life. *J Palliat Med* 2007;10:686–695.
24. Nishtala PS, Bagge ML, Campbell AJ et al. Potentially inappropriate medicines in a cohort of community-dwelling older people in New Zealand. *Geriatr Gerontol Int* 2014;14:89–93.
25. De Wilde S, Carey IM, Harris T et al. Trends in potentially inappropriate prescribing amongst older UK primary care patients. *Pharmacoepidemiol Drug Saf* 2007;16:658–667.
26. Safe Alternative to Potentially High Risk Medications in the Elderly: Medication Safety Committee, Pharmacy, Quality Division. 2013. GroupHealth [on-line]. Available at <https://provider.ghc.org/open/caringForOurMembers/pharmacy/elderly.pdf> Accessed March 11, 2015.
27. Blanco-Reina E, Ariza-Zafra G, Ocano-Riola R et al. 2012 American Geriatrics Society Beers criteria: Enhanced applicability for detecting potentially inappropriate medications in European older adults? A comparison with the Screening Tool of Older Person's Potentially Inappropriate Prescriptions. *J Am Geriatr Soc* 2014;62:1217–1223.
28. Cahir C, Fahey T, Teeling M et al. Potentially inappropriate prescribing and cost outcomes for older people: A national population study. *Br J Clin Pharmacol* 2010;69:543–552.
29. O'Mahony D, Gallagher P, Ryan C et al. STOPP & START criteria: A new approach to detecting potentially inappropriate prescribing in old age. *Eur Geriatr Med* 2010;1:45–51.
30. National Service Framework for Older People. 2001. UK Department of Health [on-line]. Available at https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/198033/National_Service_Framework_for_Older_People.pdf Accessed October 8, 2014.